

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY)		2. REPORT TYPE Technical Papers		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE <div>Please see attached</div>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER 1011	
				5e. TASK NUMBER 0046	
				5f. WORK UNIT NUMBER 346204	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/PRS 5 Pollux Drive Edwards AFB CA 93524-7048				8. PERFORMING ORGANIZATION REPORT	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/PRS 5 Pollux Drive Edwards AFB CA 93524-7048				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S NUMBER(S) Please see attached	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
20030116 033					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT A	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON Leilani Richardson
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (include area code) (661) 275-5015

MEMORANDUM FOR IN-HOUSE PUBLICATIONS

FROM: PROI (TI) (STINFO)

30 Apr 98

SUBJECT: Authorization for Release of Technical Information, Control Number: AFRL-PR-ED-TP-1998-093

J. Mills and J. Boatz Theoretical Determination of the Heats of Formation of Selected High-Energy Molecules"

HEDM Conference Presentation

(Statement A)

Attachment

Theoretical Determination of the Heats of Formation of Selected High-Energy Molecules

J.D. Mills and J.A. Boatz

Propulsion Sciences and Advanced Concepts Division
Air Force Research Laboratory (AFRL/PRS)
Edwards AFB, CA 93524-7680

Introduction—

As a measure of the intrinsic energy content of a substance, the standard enthalpy of formation constitutes a critical parameter in the evaluation of the likely performance of candidate rocket fuels and propellant additives. The gas-phase heats of formation, ΔH_f^\dagger , of a series of unusual strained and substituted organic molecules have been determined in a progression of semi-empirical, *ab initio* and density functional calculations and the results have also been compared with common non-computational estimates.

Methods—

- Depth of Treatment (increasing expense and presumed accuracy)
 - I. No computation
 - II. Semi-empirical Computation
 - III. Hartree-Fock or DFT Geometry Optimization
 - IV. Hartree-Fock or DFT Minimum Energy and Hessian
 - V. MP2 Energy (at HF Geometry) and HF Hessian
 - Exp. Experiment


● Type of Approach—Determination of ΔH_f^\dagger from:

A. Investigation of the Target Molecule Alone

B. An Isodesmic (Bond-Preserving) Reaction or a Higher

Analogue:

e.g.



Gas-phase enthalpy of formation at 298 K:

$$\begin{aligned} \Delta H_f^\dagger(\text{O} \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \text{O}) &\equiv \Delta H_{f, \text{gas}, 298\text{K}}(\text{O} \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \text{O}) \\ &= \Delta(\Delta H_f'^\dagger) - [\Delta E_{\text{eq}} + \Delta E_{\text{zp}} + \Delta[H_{298\text{K}} - H_{0\text{K}}]] \end{aligned}$$

With:

$$\begin{aligned}
\Delta(\Delta H_f^{\dagger}) &= \sum_{\substack{i' \text{ molecules} \\ \text{except target}}} \nu_{i'} \Delta H_f^{\dagger}(i') \\
&= 2 \Delta H_f^{\dagger}(\text{CH}_3\text{-O-CH}_3) + \Delta H_f^{\dagger}(\text{C}(\text{CH}_3)_4) - 4 \Delta H_f^{\dagger}(\text{CH}_4) \\
\Delta E_{\text{eq}} &= \sum_i \nu_i E_{\text{eq}}(i) \\
&= 2 E_{\text{eq}}(\text{CH}_3\text{-O-CH}_3) + E_{\text{eq}}(\text{C}(\text{CH}_3)_4) - 4 E_{\text{eq}}(\text{CH}_4) \\
&\quad - E_{\text{eq}}(\text{C}_6\text{H}_6) \\
\Delta E_{\text{zp}} &= \sum_i \nu_i E_{\text{zp}}(i) \\
\Delta[\text{H}_{298\text{K}} - \text{H}_{0\text{K}}] &= \sum_i \nu_i [\text{H}_{298\text{K}}(i) - \text{H}_{0\text{K}}(i)]
\end{aligned}$$

- Individual Methods

Method IA

Benson-Group (and Atom) Additivity Rules (Empirical Parameterization)

$$\Delta H_f^\dagger = \sum_j \Delta H_{f,j}$$

groups

S.W. Benson, *Thermochemical Kinetics*, Second Ed., (Wiley, New York, 1976), and works cited therein.

Method IIA

ΔH_f^\dagger Directly from Semi-Empirical Code

Method IIB

Semi-Empirical Energies and Thermodynamics

$$\Delta H_f^\dagger(i')\text{--exp.}; E_{eq}(i)\text{--semi-empir. calc.}; \Delta E_{zp}(i)\text{--scaled SE};$$

$$\Delta[\text{H}_{298\text{K}}(\text{i})-\text{H}_{0\text{K}}(\text{i})]-\text{exp. (if avail.) or scaled SE}$$

Method IIIA

Ibrahim and Schleyer—Method of Hartree-Fock Atom Equivalents
(Empirical/Theoretical Correlation)

$$\Delta\text{H}_\text{f}^\dagger = \text{E}_\text{HF} + \sum_{\text{j atoms}} \Delta\text{E}_\text{j}$$

M.R. Ibrahim and P.v.R. Schleyer, *J. Comp. Chem.*, **6**, 157
(1985); some parameters added by present author.

Method IIIB

Hartree-Fock or DFT Energies with SE Thermodynamics

$$\Delta\text{H}_\text{f}^\dagger(\text{i}')-\text{exp.}; \text{E}_\text{eq}(\text{i})-\text{HF/DFT}; \Delta\text{E}_\text{zp}(\text{i})-\text{scaled SE};$$

$$\Delta[\text{H}_{298\text{K}}(\text{i})-\text{H}_{0\text{K}}(\text{i})]-\text{exp. (if avail.) or scaled SE}$$

Method IVB

Hartree-Fock or DFT Energies and Thermodynamics

$\Delta H_f^\dagger(i')\text{--exp.}; E_{\text{eq}}(i)\text{--HF/DFT}; \Delta E_{\text{zp}}(i)\text{--scaled HF/DFT};$
 $\Delta[H_{298\text{K}}(i)\text{--}H_{0\text{K}}(i)]\text{--exp. (if avail.) or scaled HF}$

Method VB

MP2 Energies and HF Thermodynamics

$\Delta H_f^\dagger(i')\text{--exp.}; E_{\text{eq}}(i)\text{--MP2}; \Delta E_{\text{zp}}(i)\text{--scaled HF};$
 $\Delta[H_{298\text{K}}(i)\text{--}H_{0\text{K}}(i)]\text{--exp. (if avail.) or scaled HF}$

Method Exp

Experiment (or, if not available, high level theory, *e.g.*, G2 or
variant)








//

Semi-empirical and *ab initio* calculations performed using serial and parallel GAMESS on IBM RS/6000 work stations at Air Force Research Laboratory, and tens of nodes of the IBM SP computers at the Maui High Performance Computing Center and U.S. Army Engineer Waterways Experiment Station. Gaussian 94E on the Silicon Graphics/Cray Origin 2000 at the Aeronautical Systems Center, Major Shared Resource Center, was employed for the density functional calculations. The longest program step consumed around 40 node days.








(Semi-empirical calculations employed the PM3 Hamiltonian, *ab initio* calculations within the 6-31G(d) basis, and DFT, the B3LYP functional. Scale factors for vibrational energies: HF-0.89, PM3-0.97, DFT-0.96.)

Results—








ΔH_f^\dagger (kcal/mol) of Target Molecules (Series A)

Molecule	IA	IIA	IIIA
	86.6	74.0	83.8
	-8.5	-5.4	-9.1
	-26.9	-44.6	-30.2
	320.6	306.8	281.1
	-19.8	-26.1	-27.4
	354.3	316.2	308.7
	13.9	-18.1	-1.4
RMS vs. VB	11.9	13.2	19.0

Hartree-Fock ΔH_f^\ddagger (kcal/mol) of Targets (Series B)

Molecule	IIB	IIIB	IVB	VB
	82.1	82.5	75.1	76.5
	-12.2	-9.9	-9.9	-10.2
	-25.4	-27.4	-29.8	-28.8
	332.5	333.9	329.3	315.8
	-12.5	-15.7	-18.8	-34.7
	348.0	365.0	358.0	342.7
	1.5	13.7	8.4	-8.6
RMS vs. VB	11.6	15.7	11.7	

DFT ΔH_f^\ddagger (kcal/mol) of Targets (Series B)


Molecule	IIIB	IVB
	80.1	72.8
	-11.9	-11.7
	-29.9	-32.4
	329.5	323.6
	-24.5	-28.5
	357.2	347.8
	2.1	-4.0

Conclusions—



The accuracies of a series of progressively more expensive and sophisticated theoretical methods for calculating gas-phase enthalpies of formation have been evaluated in a specialized reference set. This procedure provides parameters which, when combined with information about condensation energetics, are fundamental to quantitatively model the performance of actual propellant formulations. However, in addition, this also justifies means of economically and efficiently screening large sets of, perhaps rather speculative, compounds at a relatively low level, so as to allow valuable experimental and theoretical resources to be focused upon candidates of particular promise.


Acknowledgments—

The Maui High Performance Computing Center and the U.S. Army Engineer Waterways Experiment Station provided computational resources on their IBM/SP computers through the DoD High Performance Computing Challenge Project Initiative. The Aeronautical System Center, Major Shared Resource Center provided time on their Silicon Graphics/Cray Origin 2000.



C_6H_8 (106 basis functions) D_{2h}





$C_{17}H_{24}N_{12}$ (483 bf.) S_4

